

Platelet count as a biomarker for monitoring treatment response and disease recurrence in recurrent epithelial ovarian cancer

CURRENT STATUS: UNDER REVIEW

Journal of Ovarian Research  BMC

Qinghong Hu

The First Affiliated Hospital of Zhengzhou University

Abha Hada

B.P. Koirala Institute of Health Sciences

Liping HAN

The First affiliated Hospital of Zhengzhou university

 hanliping0825@163.com *Corresponding Author*

DOI:

10.21203/rs.3.rs-23596/v1

SUBJECT AREAS

Sexual & Reproductive Medicine *Cancer Biology*

KEYWORDS

Platelet count, D-dimer, Fibrinogen, Recurrent EOC, Prognosis, Apoptosis

Abstract

Objectives

We sought to determine the impact of pretreatment plasma platelet levels, dimerized plasmin fragment (D-dimer) and fibrinogen in recurrent epithelial ovarian cancer (EOC) and the impact of platelet levels on SKOV3 cell lines growth and responsiveness to chemotherapy.

Methods

Under approval of ethical committee, we identified 104 women with recurrent EOC who underwent treatment between January 2010 and February 2015. Reviewing clinical, laboratory, and pathologic records from this retrospective cohort, we analyzed the correlation between pretreatment plasma D-dimer, fibrinogen, platelet levels and clinicopathological parameters, progression free survival (PFS) and overall survival (OS). Inco-culture experiments human ovarian cancer SKOV3 cell lines were used to test the effect of platelet levels on tumor growth and responsiveness to docetaxel.

Results

Of the 104 recurrent EOC, thrombocytosis at diagnosis and the decrease of platelet count by less than 25% after primary therapy were associated with worse median progression free survival ($P = 0.003$; $P = 0.021$) and median overall survival ($P = 0.009$; $P = 0.009$). Mean platelet levels declined at the end of primary therapy ($P < 0.001$) and rose at recurrence ($P = 0.007$). In multivariate analysis, elevated platelet levels at primary therapy and the decrease of platelet count less than 25% after primary therapy were unfavorable prognostic factor for PFS ($P = 0.022$; $P = 0.015$) and OS ($P = 0.013$; $P = 0.007$) in recurrent EOC, but elevated plasma D-dimer and fibrinogen were not. In SKOV-3 ovarian cancer cell lines, suitable concentration platelet co-culture protected against apoptosis ($P < 0.05$).

Conclusions

Platelet count during treatment could be used as a biomarker used for monitoring the disease recurrence and predicting treatment response. And platelet with suitable concentration co-culture protected against apoptosis in SKOV3 cell line, which may explain clinical observations.

Introduction

Malignant tumors are frequently accompanied by increased risk of hematological abnormalities. Elevated plasma D-dimer, fibrinogen, and platelet levels have been found in various malignancies[1] and are correlated with poor clinical outcome[2, 3], while some do not find any correlation[4, 5].

Platelets are associated with metastasis, angiogenesis, epithelial-to-mesenchymal transition and tumor cell proliferation[6]. Pretreatment thrombocytosis in EOC patients is closely associated with more malignant disease phenotype and poorer prognosis[7]. Thrombocytosis may independently predict the disease-specific survival for EOC patients. Thrombocytosis, accompanied by increasing of platelet aggregation rates, is associated with more aggressive tumor biology in EOC [8]. Some evidence indicates that tumor growth is incubated by a paraneoplastic thrombocytosis in a paracrine circuit by thrombopoietic cytokines in ovarian cancer culture models[9]. Ovarian cancer is ranked fifth of ten leading cancer types based on the estimated new cancer cases and deaths in women in the latest statistics[10]. Although there is a high rate of remission for advanced disease after the main treatment with surgery and chemotherapy, recurrence is common in patients with over 60% of advanced stage[11]. Despite the available literature regarding elevated plasma D-dimer, fibrinogen, and platelet levels at the time of primary treatment, it is unclear if these three at the time of recurrence have clinical significance. We considered whether plasma D-dimer, fibrinogen, and platelet levels could serve as biomarkers, like CA125, to monitor treatment response and to follow up during the surveillance for recurrence of the EOC. In this study, we evaluate the association between pretreatment plasma D-dimer, fibrinogen, platelet levels, survival, and other clinic-pathologic factors in women with recurrent EOC and the impact of platelet levels on SKOV3 cell lines growth and responsiveness to chemotherapy.

Methods

Patients

Under an ethical committee-approved protocol of the First Affiliated Hospital of Zhengzhou University, we identified 104 women with recurrent EOC who underwent treatment between January 2006 and February 2015 in our hospital. The inclusion criteria for this study were as follows: Patients with (1) pathological diagnosis of EOC, (2) treatment with surgical cytoreduction performed by a gynecologic oncologist in addition to taxane and/or platinum-based chemotherapy between January 2010 and February 2015 in our hospital, and (3) adequate clinical information in the medical record. Patients were excluded from the study for the following criteria: (1) did not receive primary therapy or follow-

up at the institution of record; (2)no recurrence of disease;(3)history of other malignancies, myeloproliferative disease, inflammatory disease, splenectomy, or other confounding causes of thrombocytosis, and (4) known congenital thrombophilia, ongoing anticoagulant treatment, pregnancy, or stroke or neurosurgery within 6 months.

Measurements Of Plasma D-dimer, Fibrinogen, Platelet Levels And Ca125

Clinical data collected included patient demographics, tumor characteristics, details of treatment, and outcome data. Pre-treatment plasma D-dimer, fibrinogen, platelet levels and CA125 measurements were recorded. Then platelet levels and CA125 measurements were recorded after the completion of surgery, 6 cycles of cytotoxic chemotherapy, after treatment completion and at the time of diagnosis of recurrence. In this cohort of patients, thrombocytosis was considered to be a platelet count $\geq 300,000$ cells/ μ l, hyperfibrinogenemia was defined as plasma fibrinogen level above 4 g/l, high plasma D-dimer level was considered based on the standard cutoff level of 0.3 mg/l, consistent with published criteria[3, 10]. Postoperative gross residual disease of < 1 cm was classified as microscopic. Conversely, residual disease > 1 cm gross was defined as macroscopic. The percent of change in platelet counts after treatment compared to the counts pretherapy were calculated. To reduce the effect of treatment-induced myelosuppression, the platelet count was determined 14 days after the end of the chemotherapy and the ratio of platelet count before and after treatment was used for further analysis.

Pre-clinical Analysis

Cell lines and culture condition

The derivation of the human ovarian cancer cell lines SKOV3 was purchased from American type culture collection (ATCC) cell library. The cell lines were maintained in RPMI-1640 with 10% fetal bovine serum. Cell lines were routinely genotyped to confirm identity and tested to confirm absence of Mycoplasma. Cells were maintained at 37 °C in a humidified incubator infused with 20% O₂ and 5% CO₂.

Docetaxel

Docetaxel (Aventis Pharma S.A.) was obtained from surplus clinical samples from the clinical pharmacy associated with Intravenous Dispensing Center of the first affiliated hospital of Zhengzhou

University.

Platelet Isolation For In Assays

Platelets were prepared for in vitro assays in a manner that would remove plasma contents and nucleated cells. Whole blood was drawn from clinical volunteers into a syringe pre-loaded with 1:9 v/v 3.8% sodium citrate and mixed 1:1 v/v with tyrodes buffer lacking Mg^{2+} and Ca^{2+} . Blood was centrifuged at 1000rpm for 10 minutes, at 22 °C. The platelet-rich plasma fraction was passed through a filtration column of Sepharose 2B beads (Sigma Aldrich, St Louis, MO) loaded into a 10 ml disposable plastic syringe with a 40 μ m nylon net filter at the bottom and sepharose 2B beads previously washed in acetone 1:1 v/v, followed by 0.9% NaCl 1:1 v/v, and Buffer 1:1 v/v. Platelets were counted with a hemocytometer by phase-contrast microscopy at 400 \times magnification and concentration was adjusted to 1×10^9 /ml by platelet buffer (Glucose 5.5 mmol/L, Tris, 15 mmol/L, NaCl 0.14 mmol/L, pH 7.4).

Potential effects of platelets on apoptosis and response to chemotherapy in assays

To examine potential effects of platelets on apoptosis and response to chemotherapy, we incubated cancer cells with platelets using a tissue co-culture system and observed consistent protection against apoptosis. To assess the effect of platelets on apoptosis, cells were plated in 6-well plates at 50,000 cells per plate. At 50% confluence, media was changed to serum-free for 24-hours prior to starting treatment. After serumstarvation, platelets were isolated and added to achieve a final dose of 1×10^8 platelets/mL. Docetaxel was dosed at 5 nM based on previously published IC₅₀ levels. Controls utilized an equivalent volume of the appropriate buffer. All treatments were performed in triplicate. After 72 hours of platelet and docetaxel exposure, cell viability was assessed using Annexin V-FITC and Propidium Iodide (PI) staining (BD Pharmingen TM) by flow cytometry. In vitro experiments were performed as described. Internal controls (n = 3) were performed for each experiment given the variability in baseline apoptosis rates seen between experiments in order to avoid batch error.

Platelet Density On Apoptosis And Response To Chemotherapy

To test the effect of platelet density on apoptosis and response to chemotherapy, cells were plated in 6-well plates at 50,000 cells per plate. At 50% confluence, media was changed to serum-free for 24-hours prior to starting treatment. After serumstarvation, platelets were isolated and added to achieve

a final dose of 0.5×10^8 platelets/mL, 1×10^8 platelets/mL, 2×10^8 platelets/mL, 4×10^8 platelets/mL.

Docetaxel was dosed at 5 nM based on previously published IC50 levels. Controls utilized an equivalent volume of the appropriate buffer. All treatments were performed in triplicate. After 72 hours of platelet and docetaxel exposure or not, cell viability was assessed using Annexin V-FITC and Propidium Iodide (PI) staining (BD Pharmingen™) by flow cytometry. In vitro experiments were performed as described. Internal controls (n = 3) were performed for each experiment given the variability in baseline apoptosis rates seen between experiments in order to avoid batch error.

Statistical Analyses

All the categorical variables were analyzed by chi-square test and ANOVA test for continuous variables. Kaplan-Meier curves were used to estimate the distribution of PFS and OS, and log-rank test was performed to compare the difference between the survival curves. Variables, which were identified as statistically significant in univariate analysis, were included in the multivariate survival analysis using the Cox proportional hazard model. Progression-free survival (PFS), interval to progression, was defined as the time from the conclusion of six cycles of primary therapy to the time when the clinical diagnosis of recurrence by physical exam, laboratory evaluation, and/or imaging was reported, whichever occurred first. Overall survival (OS) was defined as the time from the conclusion of six cycles of primary chemotherapy to the time of last follow-up or death. Survival times of patients who were known to be alive and who were dead as a result of other causes were censored with the last follow-up date. All statistical analyses were performed using SPSS 17.0 statistical software. *P* values < 0.05 were considered of statistical significance.

Results

Patient clinical characteristics

Of the 104 women with recurrent EOC, 77.9% had advanced stage (III or IV), 76.9% had high-grade disease, and 57.7% had thrombocytosis at the time of diagnosis. The median age was 53 years (range 37–81 years) and the mean platelet level was $299.8000 \mu\text{L}$ (range 108,000–545,000 cells/ μL) at diagnosis.

Relationship Between Clinicopathological Parameters And Pretreatment Plasma D-dimer/fibrinogen/platelet Levels

Plasma D-dimer and fibrinogen levels pretreatment were above the normal value in 29(27.9%) and 90(86.5%) patients, respectively. The incidence of thrombocytosis was 57.7% (60 /104).Here we found that elevated platelet levels were significantly associated with FIGO stage ($P = 0.021$) and histological type was associated with elevated plasma D-dimer levels ($P = 0.016$), but not with other clinicopathological parameters. (Table 1)

Table 1

Relationship between clinicopathological parameters and different levels of plasma D-dimer, fibrinogen and platelet count in 104 patients with recurrent EOC

Variable	n	Platelet count 1000cells/ul			Plasma Fibrinogen g/l			Plasma D-dimer mg/l		
		≤ 300	> 300	P	2-4	> 4	P	≤ 0.3	> 0.3	P
Age, y				0.340			0.386			0.566
≤55	39	28	21		27	12		5	34	
> 55	65	26	39		48	17		9	56	
Stage				0.335			0.473			> 0.05
I or and II	23	11	12		16	7		4	19	
III or and IV	81	33	48		59	22		10	71	
Histological type				0.432			0.016			0.402
Serous	64	28	36		41	23		6	58	
Non serous	40	16	24		34	6		8	32	
Grade				0.021			0.273			> 0.05
Low	24	15	9		19	5		4	20	
High	80	29	51		56	24		10	70	
Postoperative residual disease				0.069			0.514			0.402
Microscopic	66	32	34		48	18		8	58	
Macroscopic	38	12	26		27	11		6	32	

CI: confidence interval; HR: hazard ratio.

Factors affecting progression free survival and overall survival: by univariate and multivariate analyses

Of the 104 women with recurrent EOC, pretreatment thrombocytosis was associated with worse median progression free survival (11.6 vs.14.1months, $P = 0.003$, Fig. 1A) and median overall survival (15.6vs. 21.3months, $P = 0.009$, Fig. 1B).However, elevated plasma D-dimer and fibrinogen were not confirmed in univariate analysis and multivariate analysis (Table 2 and Table 3). Next we investigated the association between the percentage of reduction of platelet count relative to the pretreatment level and patient survival. Intriguingly, we found that a reduction less than 25% in platelet count post-treatment was associated with poor PFS and OS ($P = 0.021$, $P = 0.009$, Fig. 1C and 1D). After adjustment for tumor stage, histological grading, and residual disease after cytoreductive surgery,

multivariate analysis rendered decrease of platelets to a cut-off value of less than 25% reduction after treatment as an independent, unfavorable prognostic factor for PFS. (Table 2 and Table 3)

Table 2
Univariate and multivariate analysis of prognostic factors in progression free survival

Variable	Univariate analysis				Multivariate analysis			
	HR	95.0% CI		Pvalue	HR	95.0% CI		P value
Age (y) ≤ 55 vs. >55	1.351	0.903	-2.020	0.143	-	-	-	-
Postoperative residual disease Microscopic vs. Macroscopic	3.637	2.376	-5.566	0.000	2.761	1.752	-4.351	0.000
Grade Low vs. High	2.499	1.535	-4.067	0.000	1.912	1.141	-3.202	0.014
Stage I or II vs. III or IV	2.344	1.360	-4.039	0.002	1.592	0.883	-2.872	0.122
Platelet count (1000/ μ l) ≤ 300 vs. >300	1.820	1.214	-2.728	0.004	1.626	1.073	-2.466	0.022
Platelet count reduction > 25% vs. ≤25%	1.594	1.068	-2.378	0.023	1.651	1.100	-2.478	0.015
Plasma fibrinogen(g/l) 2-4 vs. >4	0.914	0.592	-1.411	0.685	-	-	-	-
Plasma D-dimer (mg/l) ≤0.3 vs. >0.3	0.894	0.507	-1.574	0.697	-	-	-	-

CI: confidence interval; HR: hazard ratio.

Table 3
Univariate and multivariate analysis of prognostic factors in overall survival

Variable	Univariate analysis				Multivariate analysis			
	HR	95.0% CI		P value	HR	95.0% CI		P value
Age (y) ≤ 55 vs. >55	1.606	0.844	-3.058	0.149	-	-	-	-
Postoperative residual disease Microscopic vs. Macroscopic	5.811	3.019	-11.182	0.000	4.300	2.124	-8.703	0.000
Grade Low vs. High	2.695	1.225	-5.928	0.014	1.714	0.733	-4.004	0.214
Stage I or II vs. III or IV	2.972	1.195	-7.393	0.019	1.799	0.656	-4.930	0.254
Platelet count (1000/ μ l) ≤ 300 vs. >300	2.298	1.208	-4.373	0.011	2.363	1.202	-4.645	0.013
Platelet count reduction > 25% vs. ≤25%	2.222	1.200	-4.113	0.011	2.363	1.262	-4.423	0.007
Plasma fibrinogen (g/l) 2-4 vs. >4	1.092	0.573	-2.083	0.789	-	-	-	-
Plasma D-dimer (mg/l) ≤0.3 vs. >0.3	0.712	0.315	-1.608	0.413	-	-	-	-

CI: confidence interval; HR: hazard ratio.

Platelet count and CA125 trends: during treatment and until the clinical diagnosis of recurrence

To consider platelet and CA125 trends, data of 104 patients was recorded through primary diagnosis, primary treatment and until the clinical diagnosis of recurrence. Of the patients with recurrent EOC (n = 104), 86.5% patients had a normal platelet count (mean 197,000 cells/ μ l) after primary therapy, and mean platelet counts at the diagnosis of recurrence were found to be increased to 227,000 cells/ μ l compared to that after primary therapy ($P = 0.007$, Fig. 2A). CA125 is a standard tumor marker followed in ovarian cancer to track the efficacy of primary therapy and in surveillance for recurrence. The mean CA125 level at diagnosis was 1034.8 IU/mL (normal < 35 IU/mL). In contrast, 81.4% had a normal CA125 level at the conclusion of primary therapy with a mean 75.1 IU/mL, and the mean post-treatment CA125 level was 12.8 IU/mL. At the clinical diagnosis of disease recurrence, CA125 was elevated in 80.8% of patients, with a median 323.1 IU/mL compared to that after treatment ($P < 0.001$, Fig. 2B). Among patients with a CA125 < 35 IU/mL at the time of recurrence, mean platelet levels at the diagnosis of recurrence were increased to 249.700 cells/ μ l compared to that (197.100 cells/ μ l) at

the conclusion of primary therapy ($P = 0.046$).

Normalization of platelet counts post-treatment is associated with disease response to therapy

In the patients with recurrent EOC who experienced a complete response to therapy > 6 months, 56.0% (51/91) had thrombocytosis at diagnosis, and 92.3% (84/91) of these patients had normalized platelet levels by the end of primary therapy. In the treatment resistant cohort ($n = 13$) who experienced a complete response to primary therapy that was durable for < 6 months, 76.9% (10/13) patients had thrombocytosis at beginning of primary therapy. At the end of primary therapy, only 46.2% (6/13) had normalized platelet counts. These data indicate a correlation between the normalization of platelet counts at the end of primary therapy and disease response to therapy ($P < 0.001$).

Platelets Resistance Against Chemotherapy-induced Apoptosis

Tissue co-culture with platelets demonstrated consistent protection against apoptosis with and without exposure to docetaxel. Platelet activation was evident by the aggregation of platelets within the initial hours of 37°C incubation. Incubation of SKOV3 cells with platelets in serum-free conditions reduced apoptosis 59.9% ($p = 0.008$). After incorporating docetaxel, incubation of the SKOV-3 cell line with platelets reduced apoptosis by 37.6% ($p = 0.0015$). (Fig. 3B and 3D)

We next compared the apoptotic rates of different platelet concentrations and tumor cells. To observe these changes in apoptotic rates, ovarian cancer cells were incubated with different concentrations of platelets for 72 hours in a serum free environment with and without docetaxel 5 nM. Incubation of SKOV3 cells with platelets of the dose 0.5×10^8 platelets/mL, 1×10^8 platelets/mL, 2×10^8 platelets/mL, 4×10^8 platelets/mL in serum-free conditions reduced apoptosis by 55.4% ($p = 0.01$), 59.9% ($p = 0.008$), 43.9% ($p = 0.044$), 8.3% ($p = 0.706$), respectively (Fig. 3A and 3C). After incorporating docetaxel, incubation of the SKOV3 cell with platelets of the dose 0.5×10^8 platelets/mL, 1×10^8 platelets/mL, 2×10^8 platelets/mL, 4×10^8 platelets/mL reduced apoptosis by 30.2% ($p = 0.03$), 37.6% ($p = 0.015$), 21.5% ($p = 0.098$), 17.4% ($p = 0.26$), respectively (Fig. 3B and 3D). These data suggest that platelets have an anti-apoptotic effect on SKOV3 cell, and they suggest that the anti-

apoptotic performance of platelets in vitro was comparable at the different seeding density. Since the culture of the SKOV3 cell with platelets incubated with 4×10^8 platelets/mL did not reduce a significantly greater number of apoptosis rates than that with 1×10^8 platelets/mL, the lower seeding density was preferable.

Discussion

Much of the research in recent years has examined the prognostic value of pretreatment platelet count in ovarian cancer[12, 13]. Notably, most of the studies have investigated the impact of the platelet count at the time of diagnosis. To our knowledge, this is the first and the largest study to evaluate the significance of plasma D-dimer, fibrinogen and platelet levels on prognosis of recurrent epithelial ovarian cancer during the course of therapy and at the time of recurrence. Also this is the first study to evaluate predictive significance of the percent of reduction of platelet level at the end of primary therapy on interval to progression and overall survival of recurrent epithelial ovarian cancer. And in this study, the result of SKOV-3 ovarian cancer cell lines co-culturing with suitable concentration platelets provides further evidence supporting a plausible role for platelet in aggressive ovarian cancer biology.

It is increasingly recognized that multiple biological components participate in a cooperative relationship between the platelets and malignancies. Cross-talk between platelets and tumor cells has been shown to participate in tumor hemorrhage[14], the epithelial-to-mesenchymal transition which promotes invasive behavior and extravasation of tumor cells, metastasis by shielding the tumor cells against NK cell mediated tumor lysis, as well as arrest of tumor cells for metastasis in target tissues[15–17]. Platelets have been shown to sequester angiogenesis regulators in addition to other mitogens and release these compounds from alpha-granules in a manner that modulates angiogenesis[18].

Previous studies from different patient cohorts have showed that elevated platelet is a poor prognostic factor[9, 19]. Here we have demonstrated through a cohort of patients with disease recurrence that elevated platelet count before treatment is associated with shortened interval to progression and decreased overall survival, but elevated plasma D-dimer and fibrinogen level were

not found to be significant, independent prognostic marker for recurrent epithelial ovarian cancer by univariate and multivariate analyses. Interestingly, here we found with strong evidence that the percent of reduction of platelet level at the end of primary treatment is an independent predictive factor of treatment response and the inadequate reduction of the platelet count is associated with shortened interval to progression and decreased overall survival of the recurrent epithelial ovarian cancer. The majority of patients with advanced stage ovarian cancer will develop recurrence. Consideration for the reduction of platelet level of the recurrent epithelial ovarian cancer patients may also help identify those who will benefit from targeted treatment. Targeting platelet counts variation tendency in ovarian cancer patients may represent a viable therapeutic target for the recurrent epithelial ovarian cancer.

The fact that there is fluctuation of platelet count during the course of primary treatment suggests its role in prediction of tumor recurrence and dependence of tumor progression. Moreover, it has been found that at the time of disease recurrence, an increased platelet count is present and was confirmed by our observations [9, 19, 20]. Platelet count is probably directly dependent on the amount of tumor cells [9] and bone marrow may be adversely affected by adjuvant chemotherapy as a result of which patients may be less likely to generate thrombocytosis in the setting of post-treatment. Disease recurrence may represent chemoresistance or altered cancer cells that have become more aggressive compared to the primary tumor. This may explain why platelet would increase at recurrence [21].

Chemoresistance is a common occurrence in recurrent EOC and platelets may contribute to this. Here we observe that elevated platelet counts at the end of primary therapy are associated with higher rates of relapse and lower rates of response to chemotherapy and the normalization of platelet counts is associated with the disease response to the primary therapy. Chemotherapy was found to be more effective in the context of thrombocytopenia in breast cancer model [22, 23] and platelets seemed to confer resistance to apoptosis induced by taxane chemotherapy [24, 25]. Pretreatment thrombocytosis and reduction of platelet level at the end of primary treatment correlate with worsened overall survival, suggesting that variation trend of platelet may be agnostic to the types of

therapy used. Furthermore, we demonstrated that pre-therapy platelet counts and reduction of platelet level at the end of primary treatment might be useful as a tumor marker, in parallel to CA125 levels, to monitor treatment response and during surveillance for recurrence.

The relationship between elevated platelets and malignancy signifies the potential role of antiplatelet drugs in the treatment of patients of EOC with thrombocytosis. There is evidence suggesting that anticoagulants may alter the potential of platelets to facilitate angiogenesis by decreasing the release of vascular endothelial growth factors[26]. A series of studies have indicated a visibly reduced malignancy risk in individuals treated with low-dose aspirin[27–31]. Nevertheless, these data need confirmation through further investigation. Based on our observation in patients with recurrent epithelial ovarian cancer, we hypothesized and confirmed that platelets with suitable concentration might confer resistance to apoptosis by taxane chemotherapy or not. Cancer cells migrate to the vasculature and interact with platelets resulting in tumor cell-induced platelet aggregation during haematogenous metastasis[32]. Co-incubation resulted in platelet aggregation and high concentrations of platelets were more likely to agglutinate. Here we further demonstrated that platelets of suitable concentration protected against apoptosis in co-culture with SKOV3 cell lines. Further, relative thrombocytopenia may be of therapeutic benefit, and within carefully defined safety parameters, the use of anti-platelet reagents may be considered as chemosensitizers.

This retrospective study has various limitations. Firstly, the data were limited by provider variation of plasma D-dimer, fibrinogen, platelet levels and CA125. Secondly, we did not address a mechanistic relationship between thrombocytosis and tumor apoptosis molecular biology. Lastly, standardization as well as prospective analysis could allow the development of prospective algorithms to test for the predictive value of platelet response as a biomarker for tumor response.

The majority of patients with advanced stage disease EOC will develop recurrence. As we move towards individualized cancer care based on patient specific genome analysis, patient centered outcomes will rise in importance and predictors of recurrence are more important than ever. Platelet level before pretreatment, during the therapy and at the time of recurrence may assist with the ability to predict disease progression and recurrence and may also help identify patients who will

benefit from the targeted therapy of cross-talk between platelets and tumor cells. Additionally, the introduction of the convenient prognostic factors such as platelet counts variation tendency can assist clinicians with better individualization of the therapeutic approach of recurrent EOC patient based on the risk stratification.

Declarations

Ethics approval and consent to participate:

Not Applicable.

Consent for publication:

Not Applicable.

Availability of data and materials:

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests:

The authors declare that they have no competing interests.

Funding:

This study was supported by the National Natural Science Foundation of China (No. U1604172), the Science and Technology Planning Project of Henan Province co-established by the province and the ministry (No. 201701002) and the Science and Technology Colleges Innovation Team Support Program of Henan Province (No. 18IRTSTHN024).

Acknowledgements:

Not Applicable.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Authors' contributions:

QHH is the primary author of this manuscript, designed experiments, performed platelet-related experiments, data analysis and wrote the manuscript. LPH contributed to design experiments and modified article; AH collected clinical data, translated the data from Chinese to English and polished language. All authors read and approved the final manuscript.

References

1. Gücer F, Tamussino K, Keil F, Balkanli-Kaplan P, Yüce MAJAR. Thrombocytosis in

- gynecologic malignancies. 2004;24(3b):2053–9.
2. Ilhan-Mutlu A, Starlinger P, Perkmann T, Schoppmann SF, Preusser M, Birner PJBiM. Plasma fibrinogen and blood platelet counts are associated with response to neoadjuvant therapy in esophageal cancer.9(4):327–35.
 3. Man Y-N, Wang Y-N, Hao J, Liu X, Liu C, Zhu C, et al. Pretreatment Plasma D-Dimer, Fibrinogen, and Platelet Levels Significantly Impact Prognosis in Patients With Epithelial Ovarian Cancer Independently of Venous Thromboembolism.25(1):24–32.
 4. Ouden MD, Ubachs JMH, Stoot JEGM, Wersch JWJVJEJoO, Gynecology, Biology R. Whole blood cell counts and leucocyte differentials in patients with benign or malignant ovarian tumours. 1997;72(1):73–7.
 5. Lavie O, Comerici G, Daras V, Bolger BS, Lopes A, Monaghan JMJGO. Thrombocytosis in Women with Vulvar Carcinoma.72(1):82–.
 6. Cho MS, Bottsford-Miller J, Vasquez HG, Stone R, Zand B, Kroll MH, et al. Platelets increase the proliferation of ovarian cancer cells.120(24):4869–72.
 7. Chen Y, Zhang L, Liu WX, Liu XYJAPJoCPA. Prognostic Significance of Preoperative Anemia, Leukocytosis and Thrombocytosis in Chinese Women with Epithelial Ovarian Cancer. 2015;16(3):933–9.
 8. Ma X, Wang Y, Sheng H, Tian W, Qi Z, Teng F, et al. Prognostic significance of thrombocytosis, platelet parameters and aggregation rates in epithelial ovarian cancer.40(1):178–83.
 9. RL S, AM N, IA M, J B-M FBHDH, et al. Paraneoplastic thrombocytosis in ovarian cancer. 2012;366(7):610.
 10. Siegel RL, KD M, Clinicians JAJCACJf. Cancer statistics, 2018. 2018;68(1):7.
 11. Salani R, Santillan A, Zahurak ML, Giuntoli RL, Gardner GJ, Armstrong DK, et al. Secondary cytoreductive surgery for localized, recurrent epithelial ovarian cancer:

- analysis of prognostic factors and survival outcome. 2010;109(4):685-91.
12. Menczer J, Schejter E, Geva D, Ginath S, Zakut H. Ovarian carcinoma associated thrombocytosis. Correlation with prognostic factors and with survival. 1998;19(1):82-4.
 13. Yu M, Liu L, Zhang BL, Chen Q, Niu TJAPJoCPA. Pretreatment Thrombocytosis as a Prognostic Factor in Women with Gynecologic Malignancies: a Meta-analysis. 2012;13(12):6077-81.
 14. Ho-Tin-Noe B, Goerge T, Wagner DDJCR. Platelets. Guardians of Tumor Vasculature.69(14):5623-6.
 15. Lakka Klement G, Yip TT, Cassiola F, Kikuchi L, Cervi D, Podust V, et al. Platelets actively sequester angiogenesis regulators.113(12):2835-42.
 16. Gay LJ, Felding-Habermann BJNRC. Contribution of platelets to tumour metastasis.11(2):123-34.
 17. Myriam L, Begum S. and et al. Direct Signaling between Platelets and Cancer Cells Induces an Epithelial-Mesenchymal-Like Transition and Promotes Metastasis.
 18. Battinelli EM, Markens BA, Italiano JEJB. Release of angiogenesis regulatory proteins from platelet alpha granules: modulation of physiologic and pathologic angiogenesis. 2011;118(5):1359-69.
 19. Allensworth SK, Langstraat CL, Martin JR, Lemens MA, Bakkum-Gamez JNJGO. The prognostic significance of preoperative thrombocytosis in epithelial ovarian cancer. 2013;130(3):S17-S8.
 20. Nather A, Mayerhofer K, Grimm C, Hefler L, Joura EAJAR. Thrombocytosis and anaemia in women with recurrent ovarian cancer prior to a second-line chemotherapy. 2003;23(3C):2991-4.
 21. Eggemann H, Ehrlicke J, Ignatov T, Fettke F, Semczuk A, Costa SD, et al. Platelet

Count After Chemotherapy is a Predictor for Outcome for Ovarian Cancer Patients.
Cancer Invest. 2015;33(5):193-6.

22. Peterson JE, Zurakowski D, Italiano JE, Michel LV, Connors S, Oenick M, et al. VEGF, PF4 and PDGF are elevated in platelets of colorectal cancer patients.15(2):265-73.
23. Ho-Tin-Noé B, Carbo C, Demers M, Cifuni SM, Goerge T. Pathology DDWJAJo. Innate Immune Cells Induce Hemorrhage in Tumors during Thrombocytopenia.175(4).
24. Radziwon-Balicka AMC, O'Driscoll L. Platelets increase survival of adenocarcinoma cells challenged with anticancer drugs: mechanisms and implications for chemoresistance British Journal of Pharmacology. 2012;167(4).
25. Bottsford-Miller J, Choi H-J, Dalton HJ, Stone RL, Cho MS, Haemmerle M, et al. Differential Platelet Levels Affect Response to Taxane-Based Therapy in Ovarian Cancer.21(3):602-10.
26. Battinelli EM, Markens BA, Kulenthirarajan RA, Machlus KR, Italiano JEJB. Anticoagulation inhibits tumor cell-mediated release of platelet angiogenic proteins and diminishes platelet angiogenic response. 2013;123(1):101.
27. Thun MJ, Eric J. Patrono. Carlo The role of aspirin in cancer prevention. Nature Reviews Clinical Oncology. 2012;9(5):259-67.
28. Rothwell PM, Wilson M, Price JF, Belch J, Mehta ZJL. Effect of daily aspirin on risk of cancer metastasis: a study of incident cancers during randomised controlled trials. 2012;379(9826).
29. Rothwell PM, Price JF, Fowkes FGR, Alberto Z, Maria Carla R, Gianni T, et al. Short-term effects of daily aspirin on cancer incidence, mortality, and non-vascular death: analysis of the time course of risks and benefits in 51 randomised controlled trials. 2012.
30. Platelets increase survival. of adenocarcinoma cells challenged with anticancer

drugs: mechanisms and implications for chemoresistance %J British Journal of Pharmacology.167(4).

31. Akl EA, Gunukula S, Barba M, Yosunico VED, Doormaal FFV, Kuipers S, et al. Parenteral anticoagulation in patients with cancer who have no therapeutic or prophylactic indication for anticoagulation. 2011.

32. 32.

Figures




Figure 1

Cumulative survival curves for PFS and OS according to pretreatment platelet count and the reduction of platelet count by Kaplan-Meier method in the 104 patients with recurrent EOC. A, Progression free survival for pretreatment platelet count ($P = 0.003$); B, Overall survival for pretreatment platelet count ($P = 0.009$); C, Progression free survival for the reduction of platelet count ($P = 0.021$); D, Overall survival for the reduction of platelet count ($P = 0.009$).




Figure 2

A: Platelet counts trend during the course of therapy and at the time of disease recurrence in 104 patients of recurrent EOC; B: CA125 levels trend during the course of therapy and at the time of disease recurrence in 104 patients of recurrent EOC; The platelet counts and CA125 levels are given in median with SD.




Figure 3

A: Apoptosis rates of SKOV-3 co-cultured with platelets $0.5 \times 10^8/\text{mL}$, $1 \times 10^8/\text{mL}$, $2 \times 10^8/\text{mL}$, $4 \times 10^8/\text{mL}$ compared to serum free media ;B: With the addition of 5 nM docetaxel, apoptosis rates of SKOV-3 co-cultured with platelets $0.5 \times 10^8/\text{mL}$, $1 \times 10^8/\text{mL}$, $2 \times 10^8/\text{mL}$, $4 \times 10^8/\text{mL}$ compared to only 5 nM docetaxel ;C:SKOV-3 co-cultured with platelets $0.5 \times 10^8/\text{mL}$, $1 \times 10^8/\text{mL}$, $2 \times 10^8/\text{mL}$, $4 \times 10^8/\text{mL}$, platelets decreased apoptosis from 15.7% to 7.0%(p = 0.01), 6.3%(p = 0.008), 8.8%(p = 0.044), 14.4%(p = 0.706) compared to serum free media respectively; D: With the addition of 5 nM docetaxel, platelets $0.5 \times 10^8/\text{mL}$, $1 \times 10^8/\text{mL}$, $2 \times 10^8/\text{mL}$, $4 \times 10^8/\text{mL}$ decreased apoptosis from 61.9% to 43.2%(p = 0.03), 38.6% (p = 0.015), 48.6%(p = 0.098), 51.1% (p = 0.26) compared to only 5nM docetaxel respectively.

